

Rational solutions of the Boussinesq equation

Peter Clarkson

School of Mathematics, Statistics and Actuarial Science,
University of Kent, Canterbury, CT2 7NF, UK
Email: P.A.Clarkson@kent.ac.uk

Abstract:

In this talk I shall discuss special polynomials associated with rational solutions of the Boussinesq equation, which is a soliton equation solvable by the inverse scattering method. These rational solutions, which are derived through a bilinear equation, are bounded, depend on two parameters and have an interesting structure. Further the rational solutions of the Boussinesq equation have similar appearance to rogue-wave solutions of the nonlinear Schrödinger equation studied in [1]–[5].

References:

1. P. Dubard, P. Gaillard, C. Klein and V. Matveev, On multi-rogue wave solutions of the NLS equation and positon solutions of the KdV equation, *Eur. Phys. J. Spec. Top.*, **185**, 247–258 (2010).
2. P. Dubard and V. Matveev, Multi-rogue waves solutions to the focusing NLS equation and the KP-I equation, *Nat. Hazards Earth Syst. Sci.*, **11**, 667–672 (2011).
3. P. Gaillard, Families of quasi-rational solutions of the NLS equation and multi-rogue waves, *J. Phys. A*, **44**, 435204 (2011).
4. P. Gaillard, Degenerate determinant representation of solutions of the nonlinear Schrödinger equation, higher order Peregrine breathers and multi-rogue waves, *J. Math. Phys.*, **54**, 013504 (2013).
5. Y. Ohta and J. Yang, General high-order rogue waves and their dynamics in the nonlinear Schrödinger equation, *Proc. R. Soc. Lond. Ser. A*, **468**, 1716–1740 (2012).